

## NOTES ON CENTRALITY RELATIONS, TERM CONDITIONS, AND COMMUTATORS

RALPH FREESE

Let  $\mathbf{A}$  be an algebra and let  $S$  and  $T$  be tolerances on  $\mathbf{A}$ . Let  $M(S, T)$ , or  $M^{\mathbf{A}}(S, T)$  to emphasize  $\mathbf{A}$ , be the set of all  $2 \times 2$  matrices of the form

$$(1) \quad \begin{bmatrix} p & q \\ r & s \end{bmatrix} = \begin{bmatrix} f(\mathbf{a}, \mathbf{u}) & f(\mathbf{a}, \mathbf{v}) \\ f(\mathbf{b}, \mathbf{u}) & f(\mathbf{b}, \mathbf{v}) \end{bmatrix}$$

where  $f(\mathbf{x}, \mathbf{y})$  is an  $(m+n)$ -ary polynomial of  $\mathbf{A}$ ,  $\mathbf{a} S \mathbf{b}$ , and  $\mathbf{u} T \mathbf{v}$  (componentwise, of course). The members of  $M(S, T)$  are called *S, T-matrices*.

The first exercise gives an efficient way to find  $M(S, T)$ .

### EXERCISES

1. Show that  $M(S, T)$  is the subalgebra of  $\mathbf{A}^4$  generated by

$$\left\{ \begin{bmatrix} a & a \\ b & b \end{bmatrix} : a S b \right\} \cup \left\{ \begin{bmatrix} c & d \\ c & d \end{bmatrix} : c T d \right\}$$

2. Use the symmetry of  $S$  and  $T$  to show the matrix obtained from an  $S, T$ -matrix by interchanging the rows or columns (or both) is also in  $M(S, T)$ .
3.  $M(T, T)$  is closed under taking transposes.

### CENTRALITY RELATIONS

We define four kinds of centrality, called centrality, strong centrality, weak centrality, and strong rectularity. There is a fifth centrality condition known as rectangularity which we will save for later.

Let  $\delta$  be a congruence and  $S$  and  $T$  be tolerance relations on  $\mathbf{A}$ . The above centrality relations are denoted  $\mathbf{C}(S, T; \delta)$  (centrality),  $\mathbf{S}(S, T; \delta)$  (strong centrality),  $\mathbf{W}(S, T; \delta)$  (weak centrality), and  $\mathbf{SR}(S, T; \delta)$  (strong rectularity). They hold if the appropriate implication below holds for all

$$\begin{bmatrix} p & q \\ r & s \end{bmatrix} \in M(S, T)$$

- centrality:  $p \delta q \implies r \delta s$ .
- strong rectangularity:  $p \delta s \implies r \delta s$ .
- weak centrality:  $p \delta q \delta s \implies r \delta s$ .
- strong centrality holds if both centrality and strong rectangularity hold.

Using the exercises it is easy to see that the implication defining  $\mathbf{C}(S, T; \delta)$  can be replaced by  $r \delta s \implies p \delta q$  and this is equivalent to

$$p \delta q \iff r \delta s.$$

Similar statements hold for the other conditions: weak centrality is equivalent to saying that if any three of  $p, q, r$  and  $s$  are  $\delta$  related, then they all are. And strong rectangularity says that if the elements of the main diagonal, or of the sinister diagonal, are  $\delta$  related, then all four are.

The  $S, T$ -term condition is the condition  $\mathbf{C}(S, T, 0)$ , usually expressed using the right-hand matrix in (1). Other kinds of term conditions are defined similarly.

If  $\mathbf{C}(S, T; \delta_i)$  holds for all  $i \in I$ , then  $\mathbf{C}(S, T; \bigwedge_{i \in I} \delta_i)$  holds. Similar statements hold for the other centrality conditions. So there is a least  $\delta$  such that  $\mathbf{C}(S, T; \delta)$  holds. This  $\delta$  is the *commutator* of  $S$  and  $T$ , and is denoted  $[S, T]$ . The commutators for the other centrality relations are denoted  $[S, T]_{\mathbf{S}}$ ,  $[S, T]_{\mathbf{SR}}$ , and  $[S, T]_{\mathbf{W}}$ .

The properties of these centrality relations are covered in Theorem 2.19 and Theorem 3.4 of [3]. Much stronger properties hold in congruence modular varieties; see [1].

#### EXERCISES

- As defined in [2],  $\beta$  is *strongly Abelian* over  $\delta$  ( $\delta \leq \beta$ , both congruences on  $\mathbf{A}$ ) if the following implication holds for all polynomials  $f$  and all elements  $x_0, \dots, x_{n-1}, y_0, \dots, y_{n-1}$ , and  $z_1, \dots, z_{n-1}$  with  $x_0 \beta y_0$  and  $x_i \beta y_i \beta z_i, i = 1, \dots, n-1$ .

$$\begin{aligned} f(x_0, \dots, x_{n-1}) \delta f(y_0, \dots, y_{n-1}) \\ \implies f(x_0, z_1, \dots, z_{n-1}) \delta f(y_0, z_1, \dots, z_{n-1}) \end{aligned}$$

Show that  $\beta$  is strongly Abelian over  $\delta$  if and only if  $\mathbf{S}(\beta, \beta; \delta)$  holds, and also show this is in turn equivalent to  $\mathbf{SR}(\beta, \beta; \delta)$ .

#### REFERENCES

- [1] Ralph Freese and Ralph McKenzie, *Commutator theory for congruence modular varieties*, London Mathematical Society Lecture Note Series, vol. 125, Cambridge University Press, Cambridge, 1987, Online version available at: <http://www.math.hawaii.edu/~ralph/papers.html>.
- [2] D. Hobby and R. McKenzie, *The structure of finite algebras (tame congruence theory)*, Contemporary Mathematics, American Mathematical Society, Providence, RI, 1988.
- [3] Keith A. Kearnes and Emil W. Kiss, *The shape of congruence lattices*.

(Ralph Freese) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAWAII, HONOLULU, HAWAII, 96822 USA

*E-mail address*, Ralph Freese: [ralph@math.hawaii.edu](mailto:ralph@math.hawaii.edu)