

Sets of equations implying semidistributivity and n -permutability

Ralph Freese

- Σ is a set of equations.

Definitions

- Σ is a set of equations.
- Σ is idempotent: if F is a function symbol occurring in Σ then $\Sigma \models x \approx F(x, \dots, x)$.

Definitions

- Σ is a set of equations.
- Σ is idempotent: if F is a function symbol occurring in Σ then $\Sigma \models x \approx F(x, \dots, x)$.
- A variety \mathcal{V} **realizes** Σ if the function symbols occurring in Σ can be interpreted as \mathcal{V} -terms such that the equations of Σ hold.

- Σ is a set of equations.
- Σ is idempotent: if F is a function symbol occurring in Σ then $\Sigma \models x \approx F(x, \dots, x)$.
- A variety \mathcal{V} **realizes** Σ if the function symbols occurring in Σ can be interpreted as \mathcal{V} -terms such that the equations of Σ hold.
- If $x \approx F(\mathbf{w})$, where \mathbf{w} is a vector of not necessarily distinct variables, then F is **weakly independent** of its i^{th} place for each i with $w_i \neq x$.

- Σ is a set of equations.
- Σ is idempotent: if F is a function symbol occurring in Σ then $\Sigma \models x \approx F(x, \dots, x)$.
- A variety \mathcal{V} **realizes** Σ if the function symbols occurring in Σ can be interpreted as \mathcal{V} -terms such that the equations of Σ hold.
- If $x \approx F(\mathbf{w})$, where \mathbf{w} is a vector of not necessarily distinct variables, then F is **weakly independent** of its i^{th} place for each i with $w_i \neq x$. So a Maltsev term $p(x, y, z)$ is weakly independent of all of its places.

Definitions

- Σ is a set of equations.
- Σ is idempotent: if F is a function symbol occurring in Σ then $\Sigma \models x \approx F(x, \dots, x)$.
- A variety \mathcal{V} **realizes** Σ if the function symbols occurring in Σ can be interpreted as \mathcal{V} -terms such that the equations of Σ hold.
- If $x \approx F(\mathbf{w})$, where \mathbf{w} is a vector of not necessarily distinct variables, then F is **weakly independent** of its i^{th} place for each i with $w_i \neq x$. So a Maltsev term $p(x, y, z)$ is weakly independent of all of its places.
- Σ' , the **derivative** is the augmentation of Σ by equations that say that F is independent of its i^{th} place whenever Σ implies F is weakly independent of its i^{th} place.

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

- If Σ' is inconsistent then any variety that realizes Σ is congruence modular (CM).

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

- If Σ' is inconsistent then any variety that realizes Σ is congruence modular (CM).
- If \mathcal{V} is a CM variety, then \mathcal{V} realizes some Σ such that Σ' is inconsistent. (The Day terms work.)

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

- If Σ' is inconsistent then any variety that realizes Σ is congruence modular (CM).
- If \mathcal{V} is a CM variety, then \mathcal{V} realizes some Σ such that Σ' is inconsistent. (The Day terms work.)
- The converse of the first statement is false: if Σ is the lattice axioms, then $\Sigma' = \Sigma$. But

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

- If Σ' is inconsistent then any variety that realizes Σ is congruence modular (CM).
- If \mathcal{V} is a CM variety, then \mathcal{V} realizes some Σ such that Σ' is inconsistent. (The Day terms work.)
- The converse of the first statement is false: if Σ is the lattice axioms, then $\Sigma' = \Sigma$. But
- The converse of the first statement is true if Σ is linear (no nested composition in the terms occurring in Σ).

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

- If Σ' is inconsistent then any variety that realizes Σ is congruence modular (CM).
- If \mathcal{V} is a CM variety, then \mathcal{V} realizes some Σ such that Σ' is inconsistent. (The Day terms work.)
- The converse of the first statement is false: if Σ is the lattice axioms, then $\Sigma' = \Sigma$. But
- The converse of the first statement is true if Σ is linear (no nested composition in the terms occurring in Σ).
- For a finite linear, idempotent Σ one can effectively decide if Σ implies CM.

The Theorems of Dent, Kearnes, Szendrei

Assume throughout this talk that Σ is idempotent. Then

- If Σ' is inconsistent then any variety that realizes Σ is congruence modular (CM).
- If \mathcal{V} is a CM variety, then \mathcal{V} realizes some Σ such that Σ' is inconsistent. (The Day terms work.)
- The converse of the first statement is false: if Σ is the lattice axioms, then $\Sigma' = \Sigma$. But
- The converse of the first statement is true if Σ is linear (no nested composition in the terms occurring in Σ).
- For a finite linear, idempotent Σ one can effectively decide if Σ implies CM. This contrasts **McNulty's Theorem** that there is no effective way to decide if a (nonlinear) idempotent Σ implies CM.

The Theorems of Dent, Kearnes, Szendrei

- A similar theorem holds for \mathcal{V} satisfying some congruence identity if

" Σ' is inconsistent"

is replaced by

" $\Sigma^{(k)}$ is inconsistent for some k ."

- The **order derivative**, Σ^+ , augments Σ by

$$x \approx F(\mathbf{w}')$$

whenever $\Sigma \models x \approx F(\mathbf{w})$, where \mathbf{w}' is the same as \mathbf{w} in every place except one, say i , and $\mathbf{w}'_i = x$.

- The **order derivative**, Σ^+ , augments Σ by

$$x \approx F(\mathbf{w}')$$

whenever $\Sigma \models x \approx F(\mathbf{w})$, where \mathbf{w}' is the same as \mathbf{w} in every place except one, say i , and $\mathbf{w}'_i = x$.

- If some iterated order derivative Σ^{+^k} of Σ is inconsistent then any variety that realizes Σ is congruence n -permutable, for some n .

- The **order derivative**, Σ^+ , augments Σ by

$$x \approx F(\mathbf{w}')$$

whenever $\Sigma \models x \approx F(\mathbf{w})$, where \mathbf{w}' is the same as \mathbf{w} in every place except one, say i , and $\mathbf{w}'_i = x$.

- If some iterated order derivative Σ^{+k} of Σ is inconsistent then any variety that realizes Σ is congruence n -permutable, for some n .
- If \mathcal{V} is a congruence n -permutable, for some n , then \mathcal{V} realizes some Σ whose iterated order derivative Σ^{+k} is inconsistent. (The Hagemann-Mitschke terms work.)

- The **order derivative**, Σ^+ , augments Σ by

$$x \approx F(\mathbf{w}')$$

whenever $\Sigma \models x \approx F(\mathbf{w})$, where \mathbf{w}' is the same as \mathbf{w} in every place except one, say i , and $\mathbf{w}'_i = x$.

- If some iterated order derivative Σ^{+k} of Σ is inconsistent then any variety that realizes Σ is congruence n -permutable, for some n .
- If \mathcal{V} is a congruence n -permutable, for some n , then \mathcal{V} realizes some Σ whose iterated order derivative Σ^{+k} is inconsistent. (The Hagemann-Mitschke terms work.)
- The converse of the first statement is false. But

- The **order derivative**, Σ^+ , augments Σ by

$$x \approx F(\mathbf{w}')$$

whenever $\Sigma \models x \approx F(\mathbf{w})$, where \mathbf{w}' is the same as \mathbf{w} in every place except one, say i , and $\mathbf{w}'_i = x$.

- If some iterated order derivative Σ^{+k} of Σ is inconsistent then any variety that realizes Σ is congruence n -permutable, for some n .
- If \mathcal{V} is a congruence n -permutable, for some n , then \mathcal{V} realizes some Σ whose iterated order derivative Σ^{+k} is inconsistent. (The Hagemann-Mitschke terms work.)
- The converse of the first statement is false. But
- The converse of the first statement is true if Σ is linear.

- The **order derivative**, Σ^+ , augments Σ by

$$x \approx F(\mathbf{w}')$$

whenever $\Sigma \models x \approx F(\mathbf{w})$, where \mathbf{w}' is the same as \mathbf{w} in every place except one, say i , and $\mathbf{w}'_i = x$.

- If some iterated order derivative Σ^{+k} of Σ is inconsistent then any variety that realizes Σ is congruence n -permutabile, for some n .
- If \mathcal{V} is a congruence n -permutabile, for some n , then \mathcal{V} realizes some Σ whose iterated order derivative Σ^{+k} is inconsistent. (The Hagemann-Mitschke terms work.)
- The converse of the first statement is false. But
- The converse of the first statement is true if Σ is linear.
- For a finite linear, idempotent Σ one can effectively decide if Σ implies congruence n -permutability, for some n .

Semidistributivity

- The **weak derivative**, Σ^* , augments Σ by an equation expressing that F is independent of its i^{th} place whenever

$$\Sigma \models x \approx F(x, \dots, x, y, x, \dots, x)$$

where the y is in the i^{th} place.

- The **weak derivative**, Σ^* , augments Σ by an equation expressing that F is independent of its i^{th} place whenever

$$\Sigma \models x \approx F(x, \dots, x, y, x, \dots, x)$$

where the y is in the i^{th} place.

- If some iterated weak derivative Σ^{*k} of Σ is inconsistent then any variety that realizes Σ is congruence semidistributive.

Semidistributivity

- The **weak derivative**, Σ^* , augments Σ by an equation expressing that F is independent of its i^{th} place whenever

$$\Sigma \models x \approx F(x, \dots, x, y, x, \dots, x)$$

where the y is in the i^{th} place.

- If some iterated weak derivative Σ^{*k} of Σ is inconsistent then any variety that realizes Σ is congruence semidistributive.
- If \mathcal{V} is a congruence semidistributive then \mathcal{V} realizes some Σ whose iterated weak derivative Σ^{*k} is inconsistent. (I'll show my variant of the Hobby-McKenzie-Kearnes-Kiss terms work.)

Semidistributivity

- The **weak derivative**, Σ^* , augments Σ by an equation expressing that F is independent of its i^{th} place whenever

$$\Sigma \models x \approx F(x, \dots, x, y, x, \dots, x)$$

where the y is in the i^{th} place.

- If some iterated weak derivative Σ^{*k} of Σ is inconsistent then any variety that realizes Σ is congruence semidistributive.
- If \mathcal{V} is a congruence semidistributive then \mathcal{V} realizes some Σ whose iterated weak derivative Σ^{*k} is inconsistent. (I'll show my variant of the Hobby-McKenzie-Kearnes-Kiss terms work.)
- The converse of the first statement is false, **even if Σ is linear**. Nevertheless

Semidistributivity

- The **weak derivative**, Σ^* , augments Σ by an equation expressing that F is independent of its i^{th} place whenever

$$\Sigma \models x \approx F(x, \dots, x, y, x, \dots, x)$$

where the y is in the i^{th} place.

- If some iterated weak derivative Σ^{*k} of Σ is inconsistent then any variety that realizes Σ is congruence semidistributive.
- If \mathcal{V} is a congruence semidistributive then \mathcal{V} realizes some Σ whose iterated weak derivative Σ^{*k} is inconsistent. (I'll show my variant of the Hobby-McKenzie-Kearnes-Kiss terms work.)
- The converse of the first statement is false, **even if Σ is linear**. Nevertheless
- For a finite linear, idempotent Σ one **can** effectively decide if Σ implies congruence semidistributivity.

My Maltsev Condition for Semidistributivity

- A variety is congruence semidistributive iff there are terms $d_i(x, y, z)$, $i = 0, \dots, n$, such that

$$d_0(x, y, z) \approx x \quad d_n(x, y, z) \approx z$$

and

My Maltsev Condition for Semidistributivity

- A variety is congruence semidistributive iff there are terms $d_i(x, y, z)$, $i = 0, \dots, n$, such that

$$d_0(x, y, z) \approx x \quad d_n(x, y, z) \approx z$$

and for each i two of the following three hold:

$$d_i(x, x, y) \approx d_{i+1}(x, x, y)$$

$$d_i(x, y, x) \approx d_{i+1}(x, y, x)$$

$$d_i(x, y, y) \approx d_{i+1}(x, y, y)$$

My Maltsev Condition for Semidistributivity

- A variety is congruence semidistributive iff there are terms $d_i(x, y, z)$, $i = 0, \dots, n$, such that

$$d_0(x, y, z) \approx x \quad d_n(x, y, z) \approx z$$

and for each i two of the following three hold:

$$d_i(x, x, y) \approx d_{i+1}(x, x, y)$$

$$d_i(x, y, x) \approx d_{i+1}(x, y, x)$$

$$d_i(x, y, y) \approx d_{i+1}(x, y, y)$$

- Let Σ be these equations. Assume inductively that Σ^{*k} implies $x \approx d_i(x, y, z)$. Then, using the above equations, one can show that $\Sigma^{*(k+2)}$ implies $x \approx d_{i+1}(x, y, z)$.

My Maltsev Condition for Semidistributivity

- A variety is congruence semidistributive iff there are terms $d_i(x, y, z)$, $i = 0, \dots, n$, such that

$$d_0(x, y, z) \approx x \quad d_n(x, y, z) \approx z$$

and for each i two of the following three hold:

$$d_i(x, x, y) \approx d_{i+1}(x, x, y)$$

$$d_i(x, y, x) \approx d_{i+1}(x, y, x)$$

$$d_i(x, y, y) \approx d_{i+1}(x, y, y)$$

- Let Σ be these equations. Assume inductively that Σ^{*k} implies $x \approx d_i(x, y, z)$. Then, using the above equations, one can show that $\Sigma^{*(k+2)}$ implies $x \approx d_{i+1}(x, y, z)$.
- So some iterated weak derivative implies $x \approx d_n(x, y, z) \approx z$ and so is inconsistent.

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

- *Is congruence modular.*

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

- *Is congruence modular.*
- *Satisfies a nontrivial congruence identity.*

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

- *Is congruence modular.*
- *Satisfies a nontrivial congruence identity.*
- *Is congruence n -permutable for some n .*

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

- *Is congruence modular.*
- *Satisfies a nontrivial congruence identity.*
- *Is congruence n -permutable for some n .*
- *Is congruence semidistributive.*

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

- *Is congruence modular.*
- *Satisfies a nontrivial congruence identity.*
- *Is congruence n -permutable for some n .*
- *Is congruence semidistributive.*
- *Is congruence meet-semidistributive.*

Theorem

For each property P listed below, given a finite, idempotent, linear set of equations Σ one can effectively decide if every variety that realizes Σ satisfies P .

- *Is congruence modular.*
- *Satisfies a nontrivial congruence identity.*
- *Is congruence n -permutable for some n .*
- *Is congruence semidistributive.*
- *Is congruence meet-semidistributive.*
- *Is congruence distributive.*