

Assignment 11 – Parts 1 & 2 – Math 411

- (1) (Projecting onto the column space of a matrix.) Let $A \in M_{m,n}(\mathbf{R})$, so that $W = \text{Col}(A) \leq V = \mathbf{R}^m$. We want to write down a formula for the orthogonal projection $P_W : V \rightarrow W$ in terms of A .
- (a) First off, for a general inner product space V and $S \subseteq V$, let $W = \text{Span}(S)$. Show that $v \in V$ is orthogonal to all $w \in W$ if and only if it is orthogonal to all $w \in S$.
 - (b) Now, let $V = \mathbf{R}^m$ (with the standard inner product) and $A \in M_{m,n}(\mathbf{R})$. Every vector in $\text{Col}(A)$ is of the form Ax for some $x \in \mathbf{R}^n$, so given $v \in V$, the orthogonal projection onto $\text{Col}(A)$ we seek is $\text{proj}_W v = A\hat{x}$ for some $\hat{x} \in \mathbf{R}^n$ satisfying $v - A\hat{x}$ is orthogonal to all $w \in \text{Col}(A)$. Show that $A\hat{x} = \text{proj}_W v$ if $A^T(v - A\hat{x}) = 0$. (Hint: use part (a) and the fact that $v_1 \cdot v_2 = v_1^T v_2$.)
 - (c) Suppose the columns of A are linearly independent, and explain why $A^T A$ is invertible.
 - (d) Suppose the columns of A are linearly independent, and show that $P_W = A(A^T A)^{-1} A^T$ (i.e. for all $v \in V$, we have $\text{proj}_W v = A(A^T A)^{-1} A^T v$).
- (2) The exercise explains a general application of inner product spaces. Let V be an inner product space and let W be a subspace of V . Given $v \in V$, we seek the closest vector in W to v . Accordingly, we say that $\bar{v} \in W$ is a *best approximation* to v in W if

$$\|v - \bar{v}\| \leq \|v - w\|, \text{ for all } w \in W.$$

- (a) The first step (carried out in parts (a)–(d)) is to show that \bar{v} is a best approximation to v if and only if $v - \bar{v}$ is orthogonal to all vectors in W (in other words, a best approximation is just what in class we called an orthogonal projection of v onto W). To show this, first suppose that $w \in W$ is such that $v - w$ is orthogonal to all vectors in W , and show that $\|v - w\| \leq \|v - w'\|$ for all $w' \in W$. (Hint: write $v - w'$ as $(v - w) - (w - w')$.)
- (b) Given a vector $w' \in W$, show that every vector w'' in W can be written as $w'' = w' - \tilde{w}$ with $\tilde{w} \in W$.
- (c) Suppose that $\|v - w\| \leq \|v - w'\|$ for all $w' \in W$. Show that this implies that for all $w'' \in W$, one has $2\text{Re}(\langle v - w, w'' \rangle) + \langle w'', w'' \rangle \geq 0$.

- (d) Plugging in $w'' = -\frac{\langle v - w, w - w' \rangle}{\langle w - w', w - w' \rangle}(w - w')$ into the inequality of part (c), conclude that if $\|v - w\| \leq \|v - w'\|$ for all $w' \in W$, then $\langle v - w, w - w' \rangle = 0$ and hence that $v - w$ is orthogonal to all vectors in W .
- (e) Show that if a best approximation exists, then it is unique.
- (3) Here's an example of applying the idea of a best approximation. The setup is the following. Let $m, n \geq 1$ and let $A \in M_{m,n}(\mathbf{R})$. Let $V = \mathbf{R}^m$ (equipped with the standard inner product) and let $W = \text{Col}(A) \leq V$. For $b \in V$, recall that there is a solution to $Ax = b$ if and only if $b \in \text{Col}(A)$. Sometimes, one has a vector $b \in V$ for which there is no solution to $Ax = b$, but there "should" be (like if b represents an experimental measurement, which should theoretically lie in the column space of A , but doesn't not due to the noise in the experiment). In such a case, we seek $\hat{x} \in \mathbf{R}^n$ such that $\|A\hat{x} - b\| \leq \|Ax - b\|$ for all $x \in \mathbf{R}^n$. Such a solution is called a *least squares solution* to $Ax = b$.
- (a) Let $p = \text{proj}_W b$ and let \hat{x} be a solution of $Ax = p$. Using Question (2) (or otherwise) show that \hat{x} is a least squares solution to $Ax = b$. (Hint: every element of W is of the form Ax for some $x \in \mathbf{R}^n$.)
- (b) Show that the only least squares solutions of $Ax = b$ are the solutions to $Ax = p$.
- (c) Conclude from Question (1) that the least squares solutions to $Ax = b$ are the solutions to $A^T Ax = A^T b$.
- (4) (Least squares linear fit.) Finally, let's give a concrete application. Suppose you have made m measurements of some physical quantity at times t_1, \dots, t_m and you obtained the values y_1, \dots, y_m , respectively. Suppose that physics tells you there should be a linear relation between y and t , i.e. there's a physical law that says that $y(t)$ satisfies $y(t) = \alpha t + \beta$ for some $\alpha, \beta \in \mathbf{R}$. How do you find α and β given the m data points? Well, one method is the least squares fit. In a perfect world, you would have that $y_i = \alpha t_i + \beta$ for all $i = 1, \dots, m$, so using two different times would yield α and β . Sadly, the world is far from perfect. Instead, you end up with m equations

$$y_1 = \alpha t_1 + \beta$$

$$\vdots$$

$$y_m = \alpha t_m + \beta$$

in the two unknowns α and β , and there's probably no solution at all! Let

$$b = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{pmatrix} \in \mathbf{R}^m \text{ and } A = \begin{pmatrix} t_1 & 1 \\ t_2 & 1 \\ \vdots & \vdots \\ t_m & 1 \end{pmatrix} \in M_{m,2}(\mathbf{R}).$$

The *least squares linear fit* of the data points $\{(t_1, y_1), \dots, (t_m, y_m)\}$ is $y(t) = \alpha t + \beta$, where $\hat{x} = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ is the least squares solution to $Ax = b$.

- (a) Suppose you have made three measurements $\{(0, 0), (1, 102), (4, 400)\}$. Find the least squares linear fit of this data.
- (5) (a) Show that the product of two unitary matrices is unitary.
 (b) Show that the product of two Hermitian matrices need not be Hermitian.
 (c) Show that the sum of two Hermitian matrices is Hermitian.
 (d) Show that the sum of two unitary matrices need not be unitary.
 (e) Can you find two unitary matrices whose sum is unitary? (If so, what are they?)
- (6) In this exercise, you'll show that the set of 2×2 orthogonal matrices consists exactly of the matrices

$$R_\theta = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix} \quad \text{and} \quad \bar{R}_\theta = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ \sin(\theta) & -\cos(\theta) \end{pmatrix}$$

for $0 \leq \theta < 2\pi$.

- (a) First, show that R_θ and \bar{R}_θ are orthogonal.
 (b) Show that, for all $\theta, \theta' \in [0, 2\pi)$, $R_\theta \neq \bar{R}_{\theta'}$. Also, show that when $\theta \neq \theta'$, $R_\theta \neq R_{\theta'}$ and $\bar{R}_\theta \neq \bar{R}_{\theta'}$.
 (c) If O is a 2×2 orthogonal matrix, show that its entries are all at most 1 in absolute value. Conclude that every entry can be written as $\cos(\theta)$ (or $\sin(\theta)$) for some $\theta \in [0, 2\pi)$.
 (d) Show that if the $(1, 1)$ -entry of O is $\pm \cos(\theta)$, then the $(2, 1)$ -entry is $\pm \sin(\theta)$ and the $(1, 2)$ -entry is also $\pm \sin(\theta)$.
 (e) Similarly, show that if the $(1, 2)$ -entry is $\pm \sin(\theta)$, then the $(2, 2)$ -entry is $\pm \cos(\theta)$.

- (f) Show that if the first column is $\begin{pmatrix} \cos(\theta) \\ \sin(\theta) \end{pmatrix}$, then the second column is $\pm \begin{pmatrix} -\sin(\theta) \\ \cos(\theta) \end{pmatrix}$.
- (g) Finally, show that you can always find $\theta \in [0, 2\pi)$ such that the first column is $\begin{pmatrix} \cos(\theta) \\ \sin(\theta) \end{pmatrix}$.
- (h) We know that R_θ represents a rotation by angle θ counterclockwise around the origin. Show that \overline{R}_θ is a reflection across the y -axis followed by R_θ . Thus, every orthogonal matrix represents either a rotation or a reflection followed by a rotation.